

K-HOMOLOGY AND FREDHOLM OPERATORS II: ELLIPTIC OPERATORS

PAUL F. BAUM AND ERIK VAN ERP

CONTENTS

1.	Introduction	1
2.	The triangle	2
2.1.	K -theory (with compact supports)	2
2.2.	Geometric K -homology	2
2.3.	The clutching map	3
2.4.	Analytic K -homology (with compact supports)	4
2.5.	The “Choose an operator” map	5
2.6.	The equivalence of geometric and analytic K -homology	6
3.	A family of index 1 operators	6
3.1.	Properties of α_F	7
3.2.	Hörmander’s operators	8
4.	The Thom class of $T(TM)$	8
5.	Proof of commutativity	11
	References	12

1. INTRODUCTION

In this paper we prove commutativity of the triangle

$$\begin{array}{ccc}
 & K^0(T^*X) & \\
 c \swarrow & & \searrow \text{Op} \\
 K_0^{\text{top}}(X) & \xrightarrow{\mu} & KK^0(C(X), \mathbb{C})
 \end{array}$$

for a closed smooth manifold X . This triangle was first introduced by Paul Baum and Ron Douglas in [3] to formulate the Atiyah-Singer index theorem for elliptic operators in the framework of K -homology. An elliptic operator on a closed manifold X determines an element in the *analytic* K -homology group $KK^0(C(X), \mathbb{C})$, and the solution of the index problem amounts to finding a *topological* description of this element. As proposed in [3], a convenient way to formalize this problem is to ask for the construction of an

PFB was partially supported by NSF grant DMS-0701184.
EvE was partially supported by NSF grant DMS-1100570.

explicit K -cycle in the *geometric* K -homology group $K_0^{top}(X)$ that (under the natural isomorphism μ between analytic and geometric K -homology) corresponds to the given elliptic operator.

Our renewed interest in the above triangle stems from its essential role in our solution of the index problem for the Heisenberg calculus on contact manifolds [6]. In [3] Baum and Douglas noted that commutativity of the triangle is implied by the Atiyah-Singer index theorem for *families* of elliptic operators. In this paper we give a direct proof that does not rely on the families index theorem. Such a direct proof is needed in order to extend the results of [6] to the families case and the equivariant case.

Commutativity of the triangle is tantamount to the assertion that the index problem for elliptic operators reduces to the index problem for Dirac operators. Hence, we provide a simple direct proof of reduction to the Dirac case.

In section 2, the groups and maps of the triangle are defined. For the proof of commutativity, preliminaries are given in sections 3 and 4. The proof is given in section 5.

2. THE TRIANGLE

2.1. K -theory (with compact supports). Throughout this paper K -theory, denoted $K^\bullet(Y)$ for a locally compact space Y , is Atiyah-Hirzebruch K -theory, i.e. topological K -theory with compact supports. Any element in $K^0(Y)$ is given by a triple (σ, E, F) where E, F are \mathbb{C} vector bundles on Y , and $\sigma : E \rightarrow F$ is a vector bundle map which is an isomorphism outside a compact subset of Y .

In particular, if X is a closed smooth manifold, and $P : C^\infty(E^0) \rightarrow C^\infty(E^1)$ is an elliptic (pseudo)differential operator, then the principal symbol $\sigma : \pi^*E^0 \rightarrow \pi^*E^1$ determines an element in $K^0(T^*X)$. Here $\pi : T^*X \rightarrow X$ is the projection.

2.2. Geometric K -homology. Geometric K -cycles were introduced in [2, 3]. For a detailed description of the relation between geometric K -homology and analytic K -homology, see [5]. In this section we briefly recall the definition and main features of K -cycle K -homology. The theory is defined for the category of paracompact Hausdorff topological spaces. In particular, since any CW complex is paracompact Hausdorff, K -cycle K -homology is defined for any CW complex.

Given a paracompact Hausdorff topological space X , a K -cycle for X is a triple (M, E, φ) consisting of a compact (without boundary) Spin^c manifold M , a smooth \mathbb{C} vector bundle E on M , and a continuous map $\varphi : M \rightarrow X$. The collection of all such K -cycles, subject to a certain equivalence relation, forms an abelian group under disjoint union. We denote this group by $K_*^{top}(X)$.

The equivalence relation that is imposed on the K -cycles is generated by three elementary steps:

- bordism
- direct sum-disjoint union
- vector bundle modification

The abelian group $K_*^{top}(X)$ consists of the equivalence classes of K -cycles. Addition is given by disjoint union,

$$(M_0, E_0, \varphi_0) + (M_1, E_1, \varphi_1) = (M_0 \sqcup M_1, E_0 \sqcup E_1, \varphi_0 \sqcup \varphi_1)$$

and the additive inverse of a K -cycle is obtained by reversing the Spin^c structure

$$-(M, E, \varphi) = (-M, E, \varphi).$$

The group $K_*^{top}(X)$ is $\mathbb{Z}/2$ graded by the parity of the dimension of the Spin^c manifold M . In other words, $K_0^{top}(X)$ and $K_1^{top}(X)$ consist of equivalence classes of (M, E, φ) cycles for which every connected component of M is even or odd dimensional respectively.

2.3. The clutching map. Let X be a closed C^∞ manifold, not required to be oriented or even dimensional. Consider the Spin^c manifold $\Sigma X = S(TX \times \mathbb{R})$, the unit sphere bundle of $TX \times \mathbb{R}$. TX is an almost complex manifold (see Section 4 below). Therefore $S(TX \times \mathbb{R})$ is a *stably* almost complex manifold, and therefore is a Spin^c manifold with Dirac operator $D_{\Sigma X}$. Note that the dimension of the manifold ΣX is two times the dimension of X , and so is even.

ΣX is a sphere bundle over X . $\varphi : \Sigma X \rightarrow X$ is the projection $S(TX \times \mathbb{R}) \rightarrow X$. Let $B(TX)$ be the unit ball bundle of TX and $S(TX)$ the unit sphere bundle of TX . For ΣX there is the “upper hemisphere” - “lower hemisphere” decomposition

$$\Sigma X = B(TX) \cup_{S(TX)} B(TX)$$

where the first copy of $B(TX)$ is the upper hemisphere and the second copy of $B(TX)$ is the lower hemisphere. The Spin^c structure on ΣX restricted to the upper hemisphere is the Spin^c structure of TX determined by its almost complex structure.

Let $(\sigma, \pi^* E^0, \pi^* E^1)$ be a (compactly supported) symbol on TX . We shall assume that the support of σ is contained in the interior of $B(TX)$. E_σ denotes the vector bundle on ΣX

$$E_\sigma = \pi^* E^0 \cup_\sigma \pi^* E^1$$

where $\pi^* E^0$ and $\pi^* E^1$ have been restricted to $B(TX)$ which is then identified, respectively, with the upper and lower hemispheres of ΣX . $\pi^* E^0$ and $\pi^* E^1$ are then clutched together on $S(TX)$ by the vector bundle isomorphism σ .

Since the interior of the upper hemisphere of ΣX identifies with TX , there is a push-forward map $\iota_* : K^0(TX) \rightarrow K^0(\Sigma X)$.

Lemma 2.3.1. *In $K^0(\Sigma X)$*

$$\iota_*(\sigma, \pi^* E^0, \pi^* E^1) = [E_\sigma] - [\varphi^* E^1]$$

Proof. Let $\tilde{\sigma}$ be the vector bundle map $E_\sigma \rightarrow \varphi^* E^1$ that on the lower hemisphere is the identity map of $\pi^* E^1$, and on the upper hemisphere is σ . Hence, (by definition) $\iota_*(\sigma, \pi^* E^0, \pi^* E^1) = (\tilde{\sigma}, E_\sigma, \varphi^* E^1)$. Since ΣX is compact, $(\tilde{\sigma}, E_\sigma, \varphi^* E^1) = [E_\sigma] - [\varphi^* E^1]$. \square

The clutching map

$$c : K^0(TX) \rightarrow K_0^{top}(X)$$

is

$$c(\sigma, \pi^* E^0, \pi^* E^1) = (\Sigma X, E_\sigma, \varphi)$$

Remark. In geometric K -homology, the K -cycle $(\Sigma X, \varphi^* E^1, \varphi)$ bounds, and therefore is zero in $K_0^{top}(X)$.

2.4. Analytic K -homology (with compact supports). We review analytic K -homology with *compact supports*. For analytic K -homology see [7, 8].

The category of topological spaces for which this theory is defined consists of compactly generated Hausdorff topological spaces for which every compact subset is second countable (i.e., metrizable). Any CW complex has this property.

For a space X in our category we denote

$$K_j^a(X) = \varinjlim_{\Delta \subset X} KK^j(C(\Delta), \mathbb{C})$$

The direct limit is taken over compact subsets $\Delta \subseteq X$. In the category of CW complexes one could take the limit over all finite subcomplexes Δ .

We briefly review the definition of the Kasparov K -homology groups for second countable compact Hausdorff spaces Δ . Then $C(\Delta)$ is a unital separable C^* -algebra. An *even Fredholm module* $(T, H^0, H^1, \rho_0, \rho_1)$ consists of two Hilbert spaces H^0, H^1 each equipped with a $*$ -representation $\rho_j : C(\Delta) \rightarrow \mathcal{L}(H^j)$, and a bounded linear operator $T : H^0 \rightarrow H^1$, such that for all $f \in C_0(X)$

$$(T^*T - 1)\rho_0(f), \quad (TT^* - 1)\rho_1(f), \quad T\rho_0(f) - \rho_1(f)T$$

are compact operators.

An *odd Fredholm module* (T, H, ρ) consists of a Hilbert space H equipped with a $*$ -representations $\rho : C(\Delta) \rightarrow \mathcal{L}(H)$, and a *self-adjoint* bounded linear operator $T : H \rightarrow H$, such that for all $f \in C_0(X)$

$$(T^2 - 1)\rho(f), \quad T\rho(f) - \rho(f)T$$

are compact operators.

The group $KK^0(C(\Delta), \mathbb{C})$ consists of equivalence classes of even Fredholm modules, while $KK^1(C(\Delta), \mathbb{C})$ consists of equivalence classes of odd Fredholm modules. Addition is defined by direct sum of Fredholm modules. The equivalence relation for Fredholm modules is generated by the direct sum relation and operator homotopy (for details, see [7, 8]).

2.5. The “Choose an operator” map. We review (including excision) the connection between elliptic operators trivial at infinity and compactly supported K -homology.

Let W be a (not necessarily compact) manifold without boundary. A pseudodifferential operator $P : C_c^\infty(W, E) \rightarrow C_c^\infty(W, F)$ of order zero is *trivial at infinity* if there exists open sets Ω_1, Ω_2 in W and an isomorphism ψ of vector bundles $E|_{\Omega_2} \rightarrow F|_{\Omega_2}$, such that

- Ω_1 has compact closure $\bar{\Omega}_1$.
- $W = \Omega_1 \cup \Omega_2$.
- If a section $s \in C_c^\infty(W, E)$ has support in Ω_1 then the support of Ps is in Ω_1 .
- If a section $s \in C_c^\infty(W, E)$ has support in Ω_2 then $Ps = \psi s$.

There is a well-defined “choose an operator” map

$$K^0(TW) \rightarrow KK_c^0(C_0(W), \mathbb{C})$$

even if W is not compact. Given a compactly supported symbol (σ, π^*E, π^*F) on TW , i.e., an element in $K^0(TW)$, we construct an element in compactly supported K -homology $KK_c^0(C_0(W), \mathbb{C})$ as follows.

First, we may assume that outside a compact set σ is the pull-back via π of a vector bundle isomorphism. Granted this, we can choose a pseudodifferential elliptic operator P of order zero that is trivial at infinity with symbol σ , with properties as listed above.

If Δ is an open set with smooth boundary and compact closure that contains Ω_1 , then due to the properties listed above P maps $C_c^\infty(\Delta, E)$ to $C_c^\infty(\Delta, F)$. This “restriction of P to Δ ” extends to a bounded operator

$$P_\Delta : L^2(\Delta, E) \rightarrow L^2(\Delta, F)$$

and therefore an element $[P_\Delta] \in KK^0(C(\Delta), \mathbb{C})$. Note that $C(\Delta)$ acts on the Hilbert spaces $L^2(\Delta, E)$, $L^2(\Delta, F)$ by the evident multiplication operators.

With Δ and σ fixed, a different choice of operator Q will result in operators P_Δ, Q_Δ that differ by a compact operator. With Δ fixed, a homotopy of σ can be lifted to a homotopy of operators. Finally, with σ and P fixed, if $\Delta \subset \Delta'$, then $P_{\Delta'} \in KK^0(C(\Delta'), \mathbb{C})$ is the direct sum of the push forward of $P_\Delta \in KK^0(C(\Delta), \mathbb{C})$ with a trivial element. In summary, the K -theory class of σ determines an element in $KK_c^0(C_0(W), \mathbb{C})$.

We will need the following excision lemma.

Lemma 2.5.1. *Let W be a manifold, and let U be an open subset of W . Then there is commutativity in the diagram*

$$\begin{array}{ccc} K^0(TU) & \longrightarrow & K^0(TW) \\ \downarrow & & \downarrow \\ KK_c^0(C_0(U), \mathbb{C}) & \longrightarrow & KK_c^0(C_0(W), \mathbb{C}) \end{array}$$

Proof. We may assume that the push-forward of a symbol σ from TU to TW restricts to σ on TU . Therefore, the operator we choose on W restricts to an operator on U with symbol σ , and outside of U this operator will be trivial. \square

2.6. The equivalence of geometric and analytic K -homology. Let Δ be a finite CW complex. As was proven in [5], there is a natural isomorphism

$$\mu : K_j^{\text{top}}(\Delta) \rightarrow KK^j(C(\Delta), \mathbb{C})$$

For the compactly supported theory, the isomorphism was introduced in [4], and is constructed as follows. Let $D_E = E \otimes D$ be the Dirac operator D for the Spin^c manifold M , twisted by the vector bundle E . Form the bounded elliptic pseudodifferential operator $T = D_E(1 + D_E^* D_E)^{-1/2}$, which defines an element in

$$[D_E] \in KK^j(C(M), \mathbb{C})$$

Then

$$\mu(M, E, \varphi) = \varphi_*([D_E]).$$

where φ_* is

$$\varphi_* : K_j^a(M) \rightarrow K_j^a(X)$$

Theorem 2.6.1. *If X is a CW complex, then*

$$\mu : K_j^{\text{top}}(X) \rightarrow K_j^a(X)$$

is an isomorphism.

Proof. That μ is an isomorphism for any CW complex X follows from the result for finite CW complexes of [5]. Both groups $K_j^{\text{top}}(X)$ and $K_j^a(X)$ are the direct limit over finite subcomplexes $\Delta \subset X$. \square

3. A FAMILY OF INDEX 1 OPERATORS

We review the properties of the family of elliptic operators of index 1 introduced in [1]. See also [10]. We place it in the context of KK -theory.

3.1. Properties of α_F . Let $\pi : F \rightarrow X$ be a C^∞ \mathbb{R} vector bundle on the closed C^∞ manifold X . Then (once a connection has been chosen for F) TF can be viewed as a \mathbb{C} vector bundle over the manifold TX . The projection $TF \rightarrow TX$ is the derivative of $F \rightarrow X$. The fibers of this vector bundle are of the form $TF_x = F_x \oplus F_x$, which we view as the complex vector space $F_x \otimes \mathbb{C}$.

Let α_F be a family of operators whose total space is F with base space X , where in each fiber F_x we choose an order zero pseudodifferential elliptic operator whose principal symbol, taken as an element in $K^0(TF_x)$, is the Bott element for $TF_x = F_x \otimes \mathbb{C}$. Since the Bott element of TF_x has compact support, the family α_F consists of operators that are trivial at infinity.

The family α_F has three relevant properties.

Property 1. The principal symbol of the family α_F is the Thom class of TF , viewed as a \mathbb{C} vector bundle on TX .

Property 2. The index of α_F , as a family, is a trivial \mathbb{C} line bundle on X . In KK -theory, this can be expressed as follows. The family α_F gives an element in the group

$$KK_c(C_0(F), C(X)) := \varinjlim_{\Delta \subseteq F} KK(C(\Delta), C(X))$$

where $\Delta \subseteq F$ ranges over all compact subsets of F and

$$\pi_* : KK_c(C_0(F), C(X)) \rightarrow KK(C(X), C(X))$$

maps α_F to the unit of the ring $KK(C(X), C(X))$.

Property 3. It follows from Property 2 that if ξ is any element in $KK(C(X), \mathbb{C})$, then the Kasparov product

$$\alpha_F \# \xi \in KK_c^0(C_0(F), \mathbb{C})$$

is an element in compactly supported K -homology of F with

$$\pi_*(\alpha_F \# \xi) = \xi \in KK^0(C(X), \mathbb{C})$$

See [1] for details on α_F .

Proposition 3.1.1. *There is commutativity in the diagram*

$$\begin{array}{ccc} K^0(TX) & \xrightarrow[\cong]{\text{Thom}} & K^0(TF) \\ \text{Op} \downarrow & & \downarrow \text{Op} \\ KK(C(X), \mathbb{C}) & \xleftarrow{\pi_*} & KK_c^0(C_0(F), \mathbb{C}) \end{array}$$

Proof. Let $\sigma \in K^0(TX)$ be the symbol of the elliptic operator P on X , i.e., $P = \text{Op}(\sigma)$. The Thom isomorphism $K^0(TX) \rightarrow K^0(TF)$ maps σ to $\tau \# \sigma$, where τ is the Thom class of TF as a complex vector bundle on TX . Then $\text{Op}(\tau \# \sigma) = \alpha_F \# P$ and by Property 3 above, $\pi_*(\alpha_F \# P) = P$. \square

3.2. Hörmander's operators. For clarity of exposition, we give some indication here of Hörmander's approach [10] to the construction of α_F .

Let V be a finite dimensional \mathbb{R} vector space with a given Euclidean inner product. The Clifford algebra $\text{Cliff}(V)$ acts on the complexification of the sum of the exterior powers $\Lambda_{\mathbb{C}}^* V$. For $v \in V$, $c(v)$ denotes Clifford multiplication by v

$$c(v) : \Lambda_{\mathbb{C}}^{\text{even}} V \rightarrow \Lambda_{\mathbb{C}}^{\text{odd}} V$$

Consider the elliptic operator

$$\alpha = c(v) + D, \quad D = d + d^*$$

which acts on \mathbb{C} valued differential forms on V , and maps even to odd forms. The *total* symbol of α is $c(v + i\xi)$, which is the Bott element $\beta \in K^0(TV)$. The operator α and its total symbol β are $O(V)$ equivariant,

$$\alpha \in KK_{O(V)}^0(C_0(V), \mathbb{C}) \quad \beta \in KK_{O(V)}^0(\mathbb{C}, C_0(TV))$$

Note that here (exceptionally for this paper) $KK_{O(V)}^0(C_0(V), \mathbb{C})$ is *not* compactly supported.

If F is an \mathbb{R} vector bundle on M , as above, choose a Euclidean structure for F , thus reducing its structure group to $O(n)$. Then due to the $O(n)$ equivariance of α , a family of operators in the fibers of F is determined,

$$\alpha'_F \in KK^0(C_0(F), C(M))$$

To obtain the element

$$\alpha_F \in KK_c^0(C_0(F), C(M))$$

in compactly supported KK -theory, realized as a family of elliptic operators trivial at infinity, as in [1], a perturbation must be made on the operator α . For details of this perturbation, see [10].

Remark 3.2.1. The total symbol β of the elliptic operator α on V is its *principal* symbol when α is viewed as an element in the Weyl calculus. Since β is invertible outside the point $(0, 0)$ in TV , it follows, using the Weyl calculus, that the operator α is Fredholm. Moreover, by the Weyl calculus index theorem, the index of α is the Bott number of its Weyl symbol β , which is 1. In fact, as shown in [10], the null space of α is 1 dimensional, and is spanned by the Gaussian $e^{-|v|^2/2}$ (a section of the line bundle $V \times \Lambda_{\mathbb{C}}^0 V$ on V). Hence, the families index of α'_F is a trivial line bundle on M .

For details on the Weyl calculus, see [9].

4. THE THOM CLASS OF $T(TM)$

Let X be a closed C^∞ manifold. $T(TX)$ is a complex vector bundle over TX in two different ways. Therefore there are two Thom classes τ_0, τ_1 . One is the symbol of the Dirac operator D_{TX} of TX , while the other is the symbol of the family α_{TX} .

A Riemannian metric for the closed C^∞ manifold X makes TX into an almost complex manifold as follows. Choose normal coordinates $x = (x_1, \dots, x_n)$ on X , and let $\xi = (\xi_1, \dots, \xi_n)$ be the induced linear coordinates on the fibers of TX . Then at point $x = 0$ we let

$$J \frac{\partial}{\partial x_j} = -\frac{\partial}{\partial \xi_j} \quad J \frac{\partial}{\partial \xi_j} = \frac{\partial}{\partial x_j}$$

Let π_0 be the projection $\pi_0 : T(TX) \rightarrow TX$, obtained by viewing $T(TX)$ as the tangent bundle of the manifold TX and using the standard projection of a vector to the point from which it emanates. Hence, in this way $T(TX)$ is a complex vector bundle on TX . Let τ_0 be its Thom class. Note that τ_0 is the symbol of the Dirac operator D_{TX} of the almost complex manifold TX ,¹

$$\tau_0 = \sigma(D_{TX})$$

With notation as in 3.1.1 above, let $F = TX$. Then $TF = T(TX)$ is a complex vector bundle over TX , where this time the projection $\pi_1 : T(TX) \rightarrow TX$ is the derivative of $TX \rightarrow X$. This is *not the same* as the projection $\pi_0 : T(TX) \rightarrow TX$ above. Let τ_1 be the Thom class of this complex vector bundle. The Thom class τ_1 is the symbol of the family α_{TX} .

At first glance, it appears that there are two Thom isomorphisms

$$K^0(TX) \xrightarrow{\cong} K^0(T(TX)) : \sigma \mapsto \pi_j^* \sigma \otimes \tau_j, \quad j = 0, 1$$

where τ_0, τ_1 are the two Thom classes introduced above. However, these two Thom isomorphisms are equal:

Lemma 4.0.2. *Let $[\sigma]$ be any element of $K^0(TX)$. Then in $K^0(T(TX))$*

$$[\pi_0^* \sigma \otimes \tau_0] = [\pi_1^* \sigma \otimes \tau_1]$$

Proof. Let $\pi_X : TX \rightarrow X$ be the usual projection, and let $\pi_{TX} : T(TX) \rightarrow TX$ be the projection for the tangent bundle of TX . There is a second map from $T(TX)$ to TX , i.e., the derivative of $\pi_X : TX \rightarrow X$. Denote $\pi_0 = \pi_{TX}$ and $\pi_1 = d\pi_X$.

We can view $T(TX)$ as a fiber bundle on X via the projection $\pi_X \circ \pi_0 : T(TX) \rightarrow X$. We denote $W := T(TX)$ if we view it as a fiber bundle on X . The fiber W_x of W at a point $x \in X$ is isomorphic to a direct sum of three copies of $T_x X$,

$$W_x \cong T_x X \oplus T_x X \oplus T_x X$$

A point $w \in W_x$ will be denoted as a triple (w_0, w_h, w_v) . First, w is a tangent vector to TX . It emanates from a point $w_0 := \pi_0(w) \in T_x X$. The pair (w_h, w_v) denotes the vertical and horizontal components of the tangent vector w to the manifold TX at w_0 .

¹Let W be a Spin^c manifold, i.e., $TW \rightarrow W$ is a Spin^c vector bundle, and therefore has a Thom class τ . The symbol of the Dirac operator of W is this Thom class τ .

More precisely, the derivative of $\pi_X : TX \rightarrow X$ is a map $d\pi_X : T(TX) \rightarrow TX$, whose kernel consists of the “vertical” tangent vectors $T^{vert}TX$ to TX . There is the canonical identification $T_{w_0}^{vert}(TX) \cong T_x X$, so we have a short exact sequence

$$0 \rightarrow T_x X \rightarrow T_{w_0}(TX) \xrightarrow{d\pi_X} T_x X \rightarrow 0$$

We let $w_h := d\pi_X(w)$. Choosing a splitting of this short exact sequence we have

$$T_{w_0}(TX) \cong T_x X \oplus T_x X$$

and we let $w_v = w - w_h$.

With this notation, the projection $\pi_{TX} : T(TX) \rightarrow TX$ is the map

$$\pi_0(w_0, w_h, w_v) := w_0$$

while the derivative $d\pi_X : T(TX) \rightarrow TX$ is the map

$$\pi_1(w_0, w_h, w_v) = w_h$$

We recall the standard construction of the Thom class for a complex vector bundle $\pi : F \rightarrow B$. On F , consider the vector bundles $\pi^* \Lambda^{even} F$ and $\pi^* \Lambda^{odd} F$, the even and odd exterior powers of F . The Thom class is given by the vector bundle map

$$\tau : \pi^* \Lambda^{even} F \rightarrow \pi^* \Lambda^{odd} F$$

which at $v \in F_b$ is

$$\tau(v) := \wedge_v + \iota_v$$

where ι_v is the adjoint of $\wedge_v : \alpha \mapsto v \wedge \alpha$ for the choice of some hermitian structure on F .

The vector bundles on $T(TX)$ used to construct the two Thom classes τ_0, τ_1 are the same, i.e., the pull-back via $\pi_X \circ \pi_0 = \pi_X \circ \pi_1$ of the even and odd parts of the exterior algebra $\Lambda TX \otimes \mathbb{C}$. At a point $w = (w_0, w_h, w_v)$ in $T(TX)$, the Thom classes τ_0, τ_1 are given by

$$\tau_0(w) = \tau(w_v + iw_h) \quad w_v + iw_h \in T_x X \otimes \mathbb{C}$$

$$\tau_1(w) = \tau(w_0 + iw_v) \quad w_0 + iw_v \in T_x X \otimes \mathbb{C}$$

Let ρ be the diffeomorphism

$$\rho : T(TX) \rightarrow T(TX) : \rho(w_0, w_h, w_v) = (w_h, -w_0, w_v)$$

ρ is properly homotopic to the identity via maps ρ_t given by

$$\rho_t(w_0, w_h, w_v) := (w_0 \cos \pi t/2 + w_h \sin \pi t/2, -w_0 \sin \pi t/2 + w_h \cos \pi t/2, w_v)$$

i.e., the map

$$T(TX) \times [0, 1] \rightarrow T(TX) : (w, t) \mapsto \rho_t(w)$$

is proper.

Note that $\tau_1(w) = i\tau_0(\rho(w))$, i.e., $\tau_1 = i\rho^* \tau_0$. Also $\pi_1 = \pi_0 \circ \rho$. Therefore

$$\pi_1^* \sigma \otimes \tau_1 = i\rho^* (\pi_0^* \sigma \otimes \tau_0)$$

□

Remark 4.0.3. Consider the commutative diagram of Proposition 3.1.1 in the special case when $F = TX$,

$$\begin{array}{ccc} K^0(TX) & \xrightarrow[\cong]{\text{Thom}} & K^0(T(TX)) \\ \text{Op} \downarrow & & \downarrow \text{Op} \\ KK(C(X), \mathbb{C}) & \xleftarrow{\pi_*} & KK_c^0(C_0(TX), \mathbb{C}) \end{array}$$

Since the two Thom isomorphisms are equal, the commutativity of this diagram can be expressed as

$$[P] = \pi_*[\sigma_P \otimes D_{TX}] \quad \text{in } KK^0(C(X), \mathbb{C})$$

However, $\sigma_P \otimes D_{TX}$ has to be interpreted as an elliptic operator of order zero on TX that is trivial at infinity, and whose principal symbol is homotopic to $\sigma_P \otimes \sigma(D_{TX})$. Note that $\sigma_P \otimes \sigma(D_{TX})$ is not in any sense an operator of Dirac type.

In the next section, by passing to an appropriate compactification of TX , we will replace the somewhat non-explicit operator $\sigma_P \otimes D_{TX}$ by an actual twisted Dirac operator.

5. PROOF OF COMMUTATIVITY

In this section we give our direct proof of reduction to Dirac. The proof is direct, and does not use any form of the Atiyah-Singer index theorem.

Let P be an elliptic (pseudo)differential operator on the closed C^∞ -manifold X ,

$$P : C^\infty(X, E) \rightarrow C^\infty(X, F)$$

E and F are complex C^∞ vector bundles on X . P determines an element

$$[P] = (T, H^0, H^1, \rho_0, \rho_1) \in KK^0(C(X), \mathbb{C})$$

where

- $T = P(1 + P^*P)^{-1/2}$
- $H^0 = L^2(X, E)$ and $H^1 = L^2(X, F)$
- $\rho_j : C(X) \rightarrow \mathcal{L}(H^j)$ is $f \mapsto \mathcal{M}_f$ where \mathcal{M}_f is the multiplication operator $(\mathcal{M}_f u)(p) = f(p)u(p)$.

The Dirac operator $D_{\Sigma X}$ of the Spin^c manifold ΣX , twisted by the vector bundle E_σ , determines an element $[E_\sigma \otimes D_{\Sigma X}] \in KK^0(C(\Sigma X), \mathbb{C})$, which pushes forward to

$$\varphi_*[E_\sigma \otimes D_{\Sigma X}] \in KK^0(C(X), \mathbb{C})$$

Theorem 5.0.4. *In $KK^0(C(X), \mathbb{C})$*

$$[P] = \varphi_*[E_\sigma \otimes D_{\Sigma X}]$$

i.e., there is commutativity in the diagram

$$\begin{array}{ccc}
 & K^0(TX) & \\
 c \swarrow & & \searrow \text{Op} \\
 K_0^{\text{top}}(X) & \xrightarrow{\mu} & KK^0(C(X), \mathbb{C})
 \end{array}$$

Proof. Consider the diagram

$$\begin{array}{ccccc}
 K^0(TX) & \xrightarrow[\cong]{\text{Thom}} & K^0(T(TX)) & \xrightarrow{\iota_*} & K^0(T(\Sigma X)) \\
 \text{Op} \downarrow & & \downarrow \text{Op} & & \downarrow \text{Op} \\
 KK(C(X), \mathbb{C}) & \xleftarrow[\cong]{\pi_*} & KK_c^0(C_0(TX), \mathbb{C}) & \xrightarrow{\iota_*} & KK(C(\Sigma X), \mathbb{C}) \\
 & & \searrow \varphi_* & &
 \end{array}$$

The diagram commutes by Proposition 3.1.1 (for the square on the left), Lemma 2.5.1 (for the square on the right), and the fact that $\pi = \varphi \circ \iota$ (for the bottom row).

Starting in the upper left corner of the diagram, let (σ, π^*E, π^*F) be the symbol of the elliptic operator P . Applying the Thom isomorphism, we obtain $\tau_1 \otimes \pi_1^* \sigma$ (where τ_1 is the symbol of the family α_{TX}). By Proposition 4.0.2 this is equal to $\tau_0 \otimes \pi_0^* \sigma$ in $K^0(T(TX))$, where now τ_0 is the symbol of the Dirac operator D_{TX} of the almost complex manifold TX .

Since D_{TX} is the restriction of the Dirac operator of ΣX to the upper hemisphere, Lemma 2.3.1 implies that

$$\iota_*(\tau_0 \otimes \pi_0^* \sigma) = \sigma(D_{\Sigma X}) \otimes (E_\sigma - \varphi^* F)$$

as elements in $K^0(T(\Sigma X))$.

Commutativity of the diagram now implies that in $KK(C(X), \mathbb{C})$

$$[P] = \varphi_*(D_{\Sigma X} \otimes E_\sigma) - \varphi_*(D_{\Sigma X}) \otimes F$$

Finally, $\varphi_*(D_{\Sigma X}) = 0$ in $KK(C(X), \mathbb{C})$ because ΣX is the boundary of the ball bundle $B(TX \times \mathbb{R})$ as a Spin^c manifold. \square

REFERENCES

- [1] M. F. Atiyah and I. M. Singer, *The index of elliptic operators. I*, Ann. of Math. (2) **87** (1968), 484–530.
- [2] P. Baum and R. G. Douglas, *Index theory, bordism, and K-homology*, Operator algebras and K-theory (San Francisco, Calif., 1981), 1982, pp. 1–31.
- [3] ———, *K homology and index theory*, Operator algebras and applications, Part I (Kingston, Ont., 1980), 1982, pp. 117–173.
- [4] ———, *Relative K homology and C* algebras*, K-Theory **5** (1991), no. 1, 1–46. MR1141333
- [5] P. Baum, N. Higson, and T. Schick, *On the equivalence of geometric and analytic K-homology*, Pure Appl. Math. Q. **3** (2007), no. 1, part 3, 1–24. MR2330153
- [6] P. F. Baum and E. van Erp, *K-homology and index theory on contact manifolds*, Acta Math. **213** (2014), no. 1, 1–48. MR3261009

- [7] B. Blackadar, *K-theory for operator algebras*, Second, Mathematical Sciences Research Institute Publications, vol. 5, Cambridge University Press, Cambridge, 1998.
- [8] N. Higson and J. Roe, *Analytic K-homology*, Oxford Mathematical Monographs, Oxford University Press, Oxford, 2000. Oxford Science Publications. MR1817560
- [9] L. Hörmander, *The Weyl calculus of pseudodifferential operators*, Comm. Pure Appl. Math. **32** (1979), no. 3, 360–444. MR517939
- [10] L. Hörmander, *On the index of pseudodifferential operators*, Elliptische Differentialgleichungen, Band II, 1971, pp. 127–146. Schriftenreihe Inst. Math. Deutsch. Akad. Wissensch. Berlin, Reihe A, Heft 8.

THE PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA, 16802, USA
E-mail address: baum@math.psu.edu

DARTMOUTH COLLEGE, 6188, KEMENY HALL, HANOVER, NEW HAMPSHIRE, 03755, USA
E-mail address: jhamvanerp@gmail.com